

# Urban Scale Modeling- Houston Case Study

---

**Joe Touma**

**U. S. EPA**

**Office of air Quality Planning & Standards  
(OAQPS)**

**Research Triangle Park, NC**

**Touma.joe@epa.gov**



# Overview

---

- **Background**
- **Objectives**
- **History**
- **Methods**
  - **Emissions**
  - **Ambient concentrations**
- **Results**
- **Conclusions**

# Background

---

- **EPA recently conducted National Scale Air Toxics Assessment (NATA)**
  - **Modeled ambient concentrations, exposure & risk of 33 air toxic pollutants**
  - **Continental US modeling domain (+PR,VI)**
  - **Geographic resolution: census tract level**
  - **Shows contribution of:**
    - **major, area & other, onroad and nonroad sources; background**
- **Results on [www.epa.gov/ttn/atw/nata](http://www.epa.gov/ttn/atw/nata)**

# Urban Scale Study Objectives

---

- **Refined dispersion model analysis of urban area improving upon NATA assessment**
- **Provide improved modeling tools to better describe air quality on an urban scale**
- **Obtain a higher degree of resolution than national scale study**
- **Help identify data gaps**

# History of Urban Study

---

- **“Conduct urban-scale assessments for a number of cities” is a component of the Integrated Urban Strategy (64FR137, July 99)**
- **Published: Dispersion Modeling of Toxic Pollutants in Urban Areas, Guidance, Methodology and Applications, EPA-454/R-99-021**
- **Published: A Simplified Approach for Estimating Secondary Production of Hazardous Air Pollutants Using the OZIPR Model, EPA-454/R-99-054**

# Methods

---

- **Emission Inventory- 1996 NTI**
  - **point: correct source locations; get site specific locations for airports, landfills**
  - **non point: allocate county level emissions to 1 km grid cells**
  - **on road mobile: allocate emissions to actual road segments using local traffic counts, road locations and emission factors**
  - **non road mobile: allocate county level emissions to 1 km grid cells**

## Methods (cont.)

---

- **Air Quality Model:**
  - **ISCST3: better area source representation; includes wet/dry deposition; added simple chemical transformation**
- **Model Performance Evaluation**
  - **Comparisons with seven ambient benzene monitors**

# Results

---

- **More detailed analysis provides more realistic patterns**
- **Better agreement with monitoring data**
- **Found large concentrations (hot spots) that were not detected in the national scale analysis**
- **Assessment still cannot be used to determine fence line concentrations**



# Conclusion

---

- **There are many air quality models applicable to urban areas**
- **We illustrated the use of the ISCST3 model**
- **We can estimate concentrations for many air toxic pollutants at a census tract resolution for input into exposure & risk models**
- **Taking more care in placing the emissions will improve results**